

## High Speed Separation of Catechins by Extreme Liquid Chromatography (UHPLC) and its Application to the Analysis of Green Tea

### Introduction

The main components of catechins in green tea include epicatechin (EC), epigallocatechin (EGC, hydroxy derivative of EC), epicatechin gallate (ECg, gallic acid ester of EC), and epigallocatechin gallate (EGCg, gallic acid ester of EGC).

We examined the applicability of X-LC for ultra-high speed analysis of 7 components: the 4 catechins listed above, catechin (C), catechin gallate (Cg), and caffeine.

### Experimental

The system utilized in this experiment was a JASCO X-LC system consisting of two 3185-PU pumps, a 3080-DG degasser, a 3180-MX mixing unit, a 3067-CO column oven, a 3070-UV UV/Vis detector, a 3059-AS autosampler, and ChromNAV chromatography data system.

A commercially-available tea beverage was filtered with 0.2  $\mu\text{m}$  membrane filter. A portion of the filtrate was injected.



Jasco UHPLC System


**JASCO INC.**

28600 Mary's Court, Easton, MD 21601 USA  
Tel: (800) 333-5272, Fax: (410) 822-7526

Application Library: <http://www.jascoinc.com/applications>


## Results and Discussion

Figure 1 shows an X-LC chromatogram of a standard mixture of 7 catechins. These components were clearly separated within 5 minutes by using a gradient elution method.



**Figure 1.** X-LC chromatogram of standard mixture. Peaks: 1 = Epigallocatechin (500 µg/mL), 2 = Caffeine (10 µg/mL), 3 = Catechin (100 µg/mL), 4 = Epicatechin (100 µg/mL), 5 = Epigallocatechin gallate (25 µg/mL), 6 = Epicatechin gallate (20 µg/mL), 7 = Catechin gallate (100 µg/mL) Conditions: Column = X-PressPak AQ-C18W (2 mmID x 50 mmL, 2 µm), Column temperature = 30°C, Mobile phase: A = 0.2% phosphoric acid / acetonitrile (90/10), B = 0.2% phosphoric acid /acetonitrile (50/50), 0 min = A/B(100/0) – 0.5 min = A/B(100/0) – 4.0 min = A/B(70/30) – 5.0 min = A/B(70/30) – 5.05 min = A/B(100/0), Flow rate = 0.7 mL/min, Detection wavelength = 280 nm, Injection volume = 1 µL.

Figure 2 shows an X-LC chromatogram of the tea sample. Each component is well separated from unknown peaks.



**Figure 2.** X-LC chromatogram of commercially-available tea beverage. Conditions: the same as in the Figure 1.